

Local coordination in MgAl₂O₄:Cr nanocrystals identified by the electron paramagnetic resonance and luminescence spectroscopy



K. Lamonova¹, I. Danilenko¹, S. Orel¹, Yu. Pashkevich¹, Yu. Kazarinov², A. Prokhorov³


¹ O. O. Galkin Donetsk Institute for Physics and Engineering of NAS of Ukraine, Kyiv, 03039, Ukraine

² V. N. Karazin Kharkiv National University, Kharkiv, 61022, Ukraine; NSC Kharkiv Institute of Physics and Technology, Kharkiv, 61108, Ukraine

³ Institute of Physics of the CAS, 182 00, Prague, Czech Republic

Introduction

The improvement of optical properties of spinel-based nanostructured phosphors in comparison with single crystals can be expected [1]. The physical nature of this phenomenon is not clear until now. Since the production of single crystals is much more expensive than the making of nanostructured samples, studying the nanoparticle size effect on the electronic structure of activation centers is a reasonable problem for technological applications in optics-related fields.



MgAl₂O₄:Cr

Experimental methods

Synthesis of nanocrystals: an inverse co-precipitation method

Characterization: X-ray phase analysis; Transmission Electron Microscopy (TEM); X-ray fluorescence spectroscopy;

Local coordination study: Electron Paramagnetic Resonance (EPR); Luminescence spectroscopy

Calculation methods


Modified Crystal Field Theory (MCFT)

The method represents a new original semi-empirical approach to calculations of the electronic structure of paramagnetic ions in coordination complexes with arbitrary symmetry and numbers of ligands.

Results


Lattice parameter, Å	
Nano 700°C	Nano 1000°C
8.08133	8.06685
Average nanoparticle size, nm	
6	15
Cr ³⁺ content, wt.%	
0.0010	0.0082

TEM images of MgAl₂O₄:Cr




MgAl₂O₄:Cr nanopowders were obtained by the method of reverse co-precipitation and annealed in the air for 2 hours at 700°C and 1000°C.


PL emission spectra of MgAl₂O₄:Cr




Energy levels of a [CrO₆] coordination complex




EPR spectra of MgAl₂O₄:Cr synthesized at 700 °C




G-factor temperature dependence



EPR spectra of MgAl₂O₄:Cr synthesized at 1000 °C



Energy levels of a [CrO₆] coordination complex



Conclusion

The local structure of the Cr activation centers was studied by luminescent and EPR spectroscopy. It was observed that the most intensive luminescence signal originates from the 1000°C -sample while the EPR signal reduces to almost zero. The MCFT calculations of the Cr electron structure revealed that the luminescent signal originates from the [CrO₆] coordination complexes that have undergone the Jahn-Teller distortions, which enhance under temperature increasing. The observed EPR signals are caused by slightly distorted [CrO₆] coordination complexes and [CrO₅Vo] complexes with vacancies.

Acknowledgements

The work reported has been done in frame of the project No. 53/19-H of the NAS of Ukraine Program "Fundamental issues of creation of new nanomaterials and nanotechnologies".

The EPR measurements were carried out within infrastructure SAFMAT in FZU CAS. This research was supported by the Czech Science Foundation (GACR), Project No. 20-21069S. We acknowledge the Operational Program Research, Development and Education financed by European Structural and Investment Funds and the Czech Ministry of Education, Youth and Sports Project SOLID21 CZ.02.1.01/0.0/0.0/16_019/0000760.

Contact information

K.V.Lamonova (k.v.lamonova@ukr.net); I. Danilenko (igord69@ukr.net); S. Orel (vultur@ukr.net); Yu. Pashkevich (yu.pashkevich@gmail.com); Yu. Kazarinov (yu.kazarinov@karazin.ua); A.Prokhorov (prokhorov@fzu.cz)

