

Optical absorption in core-shell quantum antidot with donor impurity under applied co-directed electric and magnetic fields

Holovatsky V.A., <u>Chubrei M.V.</u>

Depatment of theoretical physics and computing modelling, Chernivtsi National University after Yuriy Fed'kovych.

Kotsyubynsky, 2, Chernivtsi - 58012, Ukraine.

E-mail: <u>chubrei.maryna@chnu.edu.ua</u>

(1)

INTRODUCTION

Recently, nanostructures have attracted much attention of scientists due to modern technology of their growth and wide range of their applications in electronic and optoelectronic devices. Today it has become possible to create multilayers spherical quantum dots (MSQD), which are called core-shell, core-shell-shell and others. Based on them, biosensors, fluorescent labels, light emitting devices, solar cells are already being created, and elements for the latest computer technology are being developed [1-2].

Investigation of the electronic energy spectrum, the absorption coefficient and the influence of external fields are important for the creation of new devices for modern nanoelectronics. Therefore, to select the optimal parameters of quantum dot with the required energy spectrum of radiation, and, accordingly, the preknown characteristics of devices based on them, it is necessary to perform theoretical studies of the external fields influence on energy in spherical nanostructures.

THEORETHICAL FRAMEWORK

The semiconductor MSQD Al_{0.3}Ga_{0.7}As/GaAs/Al_{0.3}Ga_{0.7}As with impurity which is placed in center of quantum well $r_i = (r_0 + r_1)/2$ is under research.

In order to investigate the external fields effect on the electron energy spectrum and wave functions in the nanosystem with impurity it is necessary to solve the Schrodinger equation $H\psi_{im}(\vec{r}) = E_{im}\psi_{im}(\vec{r})$

 $H = \left(\vec{p} - \frac{e}{c}\vec{A}\right) \frac{1}{2u(2)} \left(\vec{p} - \frac{e}{c}\vec{A}\right) + V_F(r,\theta) + V^P(r) + U(r)$

with the Hamiltonian

where is the vector potential (ris^{θ}) the electric field potektial; is the self-polarization potential, U(r) is the confining potential.

In order to solve the equation (1), the wave functions are expanded eventies of exact functions obtained without external fields [3]. (3)

and energy spectrum E_{jm} we obtain the secular equation (4) $[H_{n\ell,n'\ell} - E_{jm}\delta_{n,n'}\delta_{\ell,\ell'}] = 0$ To determine the coefficients (4)

Using the obtained energies and wave functions of an electron, the absorption coefficients can be calculated. For a spherical nanosystem, linear, third-order nonlinear and total optical absorption coefficients (OAQ) and defined as follows, respectively

$$\alpha^{(3)}(\omega) = -\omega \sqrt{\frac{\mu}{\varepsilon_R}} \frac{(5)}{\left[\left(E_{fi} - \hbar\omega\right)^2 + \left(\hbar\Gamma_{fi}\right)^2\right]} \times \left\{4\left|M_{fi}\right|^2 - \frac{\left(M_{ff} - M_{ii}\right)^2 \left[3E_{fi}^2 - 4E_{fi}^2 \hbar\omega + \hbar^2 \left(\omega^2 - \Gamma_{fi}^2\right)\right]\right]}{\left[\left(E_{fi} - \hbar\omega\right)^2 + \left(\hbar\Gamma_{fi}\right)^2\right]^2} \times \left\{4\left|M_{fi}\right|^2 - \frac{\left(M_{ff} - M_{ii}\right)^2 \left[3E_{fi}^2 - 4E_{fi}^2 \hbar\omega + \hbar^2 \left(\omega^2 - \Gamma_{fi}^2\right)\right]\right]}{\left(E_{fi}\right)^2 + \left(\hbar\Gamma_{fi}\right)^2}\right\}$$

The dipole transition matrix element is given by
$$M_{fi} = \left\langle\psi_i \left|er\cos\theta\right|\psi_f\right\rangle$$
(7)



In the present paper, we have calculated electric and magnetic field dependences of electron energy spectra and absorption coefficient from intersubband quantum transitions electron in core-shell QD with impurity using matrix method.

RESULTS AND DISCUSSION

Dependencies of electron energies (at m=0) on electric field intensity F at different values of magnetic field induction (B = 0 T, B = 15 T and B = 30T) in the Al_{0.3}Ga_{0.7}As/GaAs/Al_{0.3}Ga_{0.7}As MSQD with core radius $r_0=5nm$ (a) $r_0=15nm$ (b) are presented in Fig. 1. It can be seen that with increasing of the quantum dot core radius the value of energy decreases, instead energy values increase with increasing of the magnetic field induction, one can see the anti-crossing of energy levels (1f and 1d). It happens due to the taken-off degeneration, which is often observed in nanostructures and studied in detail in [4]..







Fig. 2. Dependence of absorption coefficient on the applied magnetic and electric fields in the Al_{0.3}Ga_{0.7}As/GaAs/Al_{0.3}Ga_{0.7}As MSQDs with r_0 =5 nm, r_1 =15 nm and r_2 =20 nm (a) and $r_0 = 15 \text{ nm}, r_1 = 25 \text{ nm} \text{ and } r_2 = 30 \text{ nm}$ (b).

Fig. 2. indicates the linear, third-order nonlinear and total OACs in the MSQD with core radius $r_0=5nm$ (a) $r_0=15nm$ (b) with impurity which is placed in the quantum well center as function of the incident photon energy. The OACs are defined by taking into account quantum transitions from ground to several excited states. Two peaks are formed by quantum transitions 1s-1p and 1p-1d correspondently. The energy distance between peaks is bigger at bigger magnetic field induction. The value of OAC decreases with the growth of the core radius, the magnetic and electric field induction.

(b) Fig. 1. Dependence of the electron energy spectra on the applied electric field at magnetic field induction B = 0 T, B = 15 T, B= 30 T in the $AI_{0.3}Ga_{0.7}As/GaAs/AI_{0.3}Ga_{0.7}As$ multilayer spherical quantum dots with $r_0 = 5$ nm, $r_1 = 15$ nm and $r_2 = 20$ nm (a) and $r_0 = 15$ nm, $r_1 = 25$ nm and $r_2 = 30$ nm (b).

CONCLUSIONS REFERENCES The dependences of electron energies and absorption coefficient on external fields intensity are 1. M.F. Frasco and N. Chaniotakis, Semiconductor quantum dots in chemical sensors and biosensors.-2009.-9.-pp. 7266-7286,. obtained. 2. X. Wang, H. Li, and G. Chen, Core-shell nanoparticles for cancer imaging and therapy. in It is shown that with increasing of the quantum dot core radius the value of energy decreases, instead Core-Shell Nanostructures for Drug Delivery and Theranostics, Elsevier, Ámsterdam.-2018.-466.-pp. energy values increase with increasing of the magnetic field induction and decrease with electric field 143–175. induction increasing. 3. Holovatsky V., Chubrey M., Voitsekhivska O. Effect of electric field on photoionisation cross-The results show that external fields and impurity presence effects cause significant changes on the section of impurity in multilayered quantum dot// Superlattices and Microstructures. -2020.- 145.optical absorption coefficient. For nanosystems with a bigger core, the external fields effect is stronger. Is P.106642 obtained that absorption coefficient peaks shift to the region of higher energies with electric fied intensity 4. A.S. Baimuratov, I.D. Rukhlenko, V.K. Turkov, I.O. Ponomareva, M.Y. Leonov, T.S. Perova, K. Berwick, A.V. Baranov, A.V. Federov, Level anticrossing of impurity states in semiconductor increasing, the energy distance between peaks is bigger at bigger magnetic field induction, for r_0 = 5 nm nanocrystals, *Scientific Reports*,-2014.- **4**.-P. 6917 linear OAC is much bigger than third-order nonlinear OAC. STU ÜLIKE

